

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Average	34.8%
	Response at Six Months	Average	8.7%
	Remission at Six Months	Average	5.8%
	Follow-up at 12 Months	Below	4.4%
	Response at 12 Months	Below	2.9%
	Remission at 12 Months	Average	1.5%
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Below	17.7%
	Response at Six Months	Average	2.9%
	Remission at Six Months	Average	0.0%
	Follow-up at 12 Months	Below	0.0%
	Response at 12 Months	Below	0.0%
	Remission at 12 Months	Average	0.0%
Adolescent Mental Health and/or Depression Screening*		Above	95.9%
Colorectal Cancer Screening		-	-
Optimal Asthma Control - Adults		Above	83.3%
Optimal Asthma Control - Children		Above	85.7%
Optimal Diabetes Care		-	-
Optimal Vascular Care		-	-

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).