

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	94.1%
	Follow-up at Six Months	Above	69.1%
	Response at Six Months	Average	16.5%
	Remission at Six Months	Average	6.6%
	Follow-up at 12 Months	Above	62.5%
	Response at 12 Months	Average	15.8%
	Remission at 12 Months	Average	6.6%
Adult Depression	PHQ-9 Utilization*	Above	96.6%
	Follow-up at Six Months	Above	78.4%
	Response at Six Months	Average	21.5%
	Remission at Six Months	Average	11.2%
	Follow-up at 12 Months	Above	77.7%
	Response at 12 Months	Average	21.6%
	Remission at 12 Months	Average	13.1%
Adolescent Mental Health and/or Depression Screening*		-	-
Colorectal Cancer Screening		-	-
Optimal Asthma Control - Adults		-	-
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		-	-
Optimal Vascular Care		-	-

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).