

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	97.5%
	Follow-up at Six Months	Average	35.5%
	Response at Six Months	Average	10.8%
	Remission at Six Months	Average	3.2%
	Follow-up at 12 Months	Average	32.3%
	Response at 12 Months	Average	7.5%
	Remission at 12 Months	Average	3.2%
Adult Depression	PHQ-9 Utilization*	Above	95.7%
	Follow-up at Six Months	Average	28.1%
	Response at Six Months	Average	9.4%
	Remission at Six Months	Average	6.3%
	Follow-up at 12 Months	Below	15.6%
	Response at 12 Months	Average	9.4%
	Remission at 12 Months	Average	9.4%
Adolescent Mental Health and/or Depression Screening*		Above	98.7%
Colorectal Cancer Screening		-	-
Optimal Asthma Control - Adults		-	-
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		-	-
Optimal Vascular Care		-	-

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).