

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	97.5%
	Follow-up at Six Months	Average	38.1%
	Response at Six Months	Average	8.9%
	Remission at Six Months	Average	4.5%
	Follow-up at 12 Months	Average	40.1%
	Response at 12 Months	Average	18.3%
	Remission at 12 Months	Average	9.9%
Adult Depression	PHQ-9 Utilization*	Above	94.4%
	Follow-up at Six Months	Below	25.6%
	Response at Six Months	Average	7.7%
	Remission at Six Months	Average	3.9%
	Follow-up at 12 Months	Below	20.5%
	Response at 12 Months	Below	5.1%
	Remission at 12 Months	Average	2.6%
Adolescent Mental Health and/or Depression Screening*		Above	96.0%
Colorectal Cancer Screening		-	-
Optimal Asthma Control - Adults		-	-
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		-	-
Optimal Vascular Care		-	-

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).