

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	35.3%
	Response at Six Months	Average	14.7%
	Remission at Six Months	Average	5.9%
	Follow-up at 12 Months	Average	32.4%
	Response at 12 Months	Average	8.8%
	Remission at 12 Months	Average	5.9%
Adult Depression	PHQ-9 Utilization*	Average	74.0%
	Follow-up at Six Months	Below	36.7%
	Response at Six Months	Below	12.6%
	Remission at Six Months	Average	6.9%
	Follow-up at 12 Months	Below	26.4%
	Response at 12 Months	Below	9.2%
	Remission at 12 Months	Below	5.4%
Adolescent Mental Health and/or Depression Screening*		Below	83.7%
Colorectal Cancer Screening		Above	71.4%
Optimal Asthma Control - Adults		Below	41.4%
Optimal Asthma Control - Children		Average	37.8%
Optimal Diabetes Care		Average	40.4%
Optimal Vascular Care		Below	33.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).