

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	48.8%
	Response at Six Months	Average	4.7%
	Remission at Six Months	Average	2.3%
	Follow-up at 12 Months	Above	60.5%
	Response at 12 Months	Average	14.0%
	Remission at 12 Months	Average	7.0%
Adult Depression	PHQ-9 Utilization*	Above	92.3%
	Follow-up at Six Months	Above	59.9%
	Response at Six Months	Above	25.1%
	Remission at Six Months	Average	14.1%
	Follow-up at 12 Months	Above	59.1%
	Response at 12 Months	Above	27.6%
	Remission at 12 Months	Average	13.8%
Adolescent Mental Health and/or Depression Screening*		Average	92.8%
Colorectal Cancer Screening		Above	75.0%
Optimal Asthma Control - Adults		Above	71.7%
Optimal Asthma Control - Children		Above	74.3%
Optimal Diabetes Care		Above	56.4%
Optimal Vascular Care		Above	65.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).