

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	56.8%
	Response at Six Months	Average	11.4%
	Remission at Six Months	Average	6.8%
	Follow-up at 12 Months	Average	45.5%
	Response at 12 Months	Average	25.0%
	Remission at 12 Months	Average	9.1%
Adult Depression	PHQ-9 Utilization*	Above	88.0%
	Follow-up at Six Months	Above	56.1%
	Response at Six Months	Above	23.1%
	Remission at Six Months	Average	12.9%
	Follow-up at 12 Months	Above	55.2%
	Response at 12 Months	Above	23.1%
	Remission at 12 Months	Average	12.5%
Adolescent Mental Health and/or Depression Screening*		Average	94.0%
Colorectal Cancer Screening		Above	70.8%
Optimal Asthma Control - Adults		Above	68.5%
Optimal Asthma Control - Children		Above	70.7%
Optimal Diabetes Care		Average	40.2%
Optimal Vascular Care		Average	58.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).