

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	82.0%
	Follow-up at Six Months	Average	47.6%
	Response at Six Months	Average	11.1%
	Remission at Six Months	Average	3.2%
	Follow-up at 12 Months	Average	27.0%
	Response at 12 Months	Average	6.4%
	Remission at 12 Months	Average	1.6%
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Above	98.7%
Colorectal Cancer Screening		-	-
Optimal Asthma Control - Adults		Average	62.5%
Optimal Asthma Control - Children		Above	62.7%
Optimal Diabetes Care		-	-
Optimal Vascular Care		-	-

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).