

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	84.3%
	Follow-up at Six Months	Average	48.0%
	Response at Six Months	Average	14.7%
	Remission at Six Months	Average	4.0%
	Follow-up at 12 Months	Average	48.0%
	Response at 12 Months	Average	10.7%
	Remission at 12 Months	Average	5.3%
Adult Depression	PHQ-9 Utilization*	Above	89.2%
	Follow-up at Six Months	Average	47.6%
	Response at Six Months	Average	18.7%
	Remission at Six Months	Average	10.5%
	Follow-up at 12 Months	Average	48.6%
	Response at 12 Months	Average	20.4%
	Remission at 12 Months	Average	13.9%
Adolescent Mental Health and/or Depression Screening*		Above	98.5%
Colorectal Cancer Screening		Above	72.6%
Optimal Asthma Control - Adults		Above	68.2%
Optimal Asthma Control - Children		Above	67.8%
Optimal Diabetes Care		Average	48.1%
Optimal Vascular Care		Average	57.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).