

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	40.4%
	Response at Six Months	Average	10.6%
	Remission at Six Months	Average	6.4%
	Follow-up at 12 Months	Average	53.2%
	Response at 12 Months	Average	12.8%
	Remission at 12 Months	Average	4.3%
Adult Depression	PHQ-9 Utilization*	Above	95.4%
	Follow-up at Six Months	Above	62.6%
	Response at Six Months	Average	25.2%
	Remission at Six Months	Average	14.4%
	Follow-up at 12 Months	Above	70.3%
	Response at 12 Months	Above	29.7%
	Remission at 12 Months	Above	17.1%
Adolescent Mental Health and/or Depression Screening*		Above	94.5%
Colorectal Cancer Screening		Average	66.8%
Optimal Asthma Control - Adults		Above	73.6%
Optimal Asthma Control - Children		Average	66.0%
Optimal Diabetes Care		Above	54.1%
Optimal Vascular Care		Average	59.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).