

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	40.2%
	Follow-up at Six Months	Average	30.0%
	Response at Six Months	Average	9.0%
	Remission at Six Months	Average	5.0%
	Follow-up at 12 Months	Below	15.0%
	Response at 12 Months	Below	6.0%
	Remission at 12 Months	Average	3.0%
Adolescent Mental Health and/or Depression Screening*		Below	51.1%
Colorectal Cancer Screening		Average	64.1%
Optimal Asthma Control - Adults		Below	9.1%
Optimal Asthma Control - Children		Below	12.3%
Optimal Diabetes Care		Average	34.5%
Optimal Vascular Care		Below	33.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).