

**Clinical Quality Measures**

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	69.6%
	Follow-up at Six Months	Average	46.0%
	Response at Six Months	Average	18.9%
	Remission at Six Months	Average	10.8%
	Follow-up at 12 Months	Average	56.8%
	Response at 12 Months	Average	27.0%
	Remission at 12 Months	Average	24.3%
<b>Adolescent Mental Health and/or Depression Screening*</b>		Below	83.3%
<b>Colorectal Cancer Screening</b>		Below	59.7%
<b>Optimal Asthma Control - Adults</b>		-	-
<b>Optimal Asthma Control - Children</b>		-	-
<b>Optimal Diabetes Care</b>		Average	42.3%
<b>Optimal Vascular Care</b>		Average	46.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

**Ratings**

**Above:** Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).