

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Average	25.0%
	Response at Six Months	Average	6.3%
	Remission at Six Months	Average	3.1%
	Follow-up at 12 Months	Average	28.1%
	Response at 12 Months	Average	9.4%
	Remission at 12 Months	Average	3.1%
Adult Depression	PHQ-9 Utilization*	Average	74.5%
	Follow-up at Six Months	Average	43.7%
	Response at Six Months	Average	20.1%
	Remission at Six Months	Average	10.1%
	Follow-up at 12 Months	Average	49.3%
	Response at 12 Months	Average	18.1%
	Remission at 12 Months	Average	10.6%
Adolescent Mental Health and/or Depression Screening*		Above	97.3%
Colorectal Cancer Screening		Above	70.5%
Optimal Asthma Control - Adults		Average	45.6%
Optimal Asthma Control - Children		Below	35.1%
Optimal Diabetes Care		Average	49.6%
Optimal Vascular Care		Average	57.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).