

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	86.4%
	Follow-up at Six Months	Average	43.6%
	Response at Six Months	Average	11.3%
	Remission at Six Months	Average	4.8%
	Follow-up at 12 Months	Average	51.6%
	Response at 12 Months	Average	25.8%
	Remission at 12 Months	Average	17.7%
Adult Depression	PHQ-9 Utilization*	Above	90.8%
	Follow-up at Six Months	Above	56.6%
	Response at Six Months	Average	20.4%
	Remission at Six Months	Average	13.5%
	Follow-up at 12 Months	Above	51.8%
	Response at 12 Months	Average	20.6%
	Remission at 12 Months	Average	10.6%
Adolescent Mental Health and/or Depression Screening*		Below	88.8%
Colorectal Cancer Screening		Average	67.0%
Optimal Asthma Control - Adults		Above	65.9%
Optimal Asthma Control - Children		Above	64.0%
Optimal Diabetes Care		Above	48.9%
Optimal Vascular Care		Above	65.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).