

Clinical Quality Measures
 2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	86.5%
	Follow-up at Six Months	Average	56.0%
	Response at Six Months	Average	23.2%
	Remission at Six Months	Average	13.2%
	Follow-up at 12 Months	Average	50.4%
	Response at 12 Months	Average	20.4%
	Remission at 12 Months	Average	13.2%
Adolescent Mental Health and/or Depression Screening*		Average	91.5%
Colorectal Cancer Screening		Below	64.1%
Optimal Asthma Control - Adults		Above	66.3%
Optimal Asthma Control - Children		Average	67.2%
Optimal Diabetes Care		Average	46.7%
Optimal Vascular Care		Average	59.3%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).