

Clinical Quality Measures

2022 measurement year

| Measure | | Rating | Rate |
|--|-------------------------|--------|-------|
| Adolescent Depression | PHQ-9 Utilization* | - | - |
| | Follow-up at Six Months | - | - |
| | Response at Six Months | - | - |
| | Remission at Six Months | - | - |
| | Follow-up at 12 Months | - | - |
| | Response at 12 Months | - | - |
| | Remission at 12 Months | - | - |
| Adult Depression | PHQ-9 Utilization* | Below | 38.8% |
| | Follow-up at Six Months | Below | 24.3% |
| | Response at Six Months | Below | 6.8% |
| | Remission at Six Months | Below | 3.9% |
| | Follow-up at 12 Months | Below | 23.3% |
| | Response at 12 Months | Below | 7.8% |
| | Remission at 12 Months | Below | 1.9% |
| Adolescent Mental Health and/or Depression Screening* | | Below | 49.9% |
| Colorectal Cancer Screening | | Above | 77.9% |
| Optimal Asthma Control - Adults | | Below | 23.4% |
| Optimal Asthma Control - Children | | Below | 15.4% |
| Optimal Diabetes Care | | Below | 39.2% |
| Optimal Vascular Care | | Below | 40.4% |

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).