

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	79.5%
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	53.4%
	Follow-up at Six Months	Below	27.5%
	Response at Six Months	Below	6.4%
	Remission at Six Months	Below	4.1%
	Follow-up at 12 Months	Below	22.8%
	Response at 12 Months	Average	9.9%
	Remission at 12 Months	Average	6.4%
Adolescent Mental Health and/or Depression Screening*		Below	75.2%
Colorectal Cancer Screening		Average	65.0%
Optimal Asthma Control - Adults		Below	22.5%
Optimal Asthma Control - Children		Below	24.6%
Optimal Diabetes Care		Average	39.5%
Optimal Vascular Care		Below	42.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).