

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	86.3%
	Follow-up at Six Months	Above	70.3%
	Response at Six Months	Average	18.8%
	Remission at Six Months	Average	10.9%
	Follow-up at 12 Months	Above	64.1%
	Response at 12 Months	Average	14.1%
	Remission at 12 Months	Average	10.9%
Adult Depression	PHQ-9 Utilization*	Above	86.4%
	Follow-up at Six Months	Above	69.2%
	Response at Six Months	Average	15.9%
	Remission at Six Months	Average	8.2%
	Follow-up at 12 Months	Above	64.7%
	Response at 12 Months	Average	17.5%
	Remission at 12 Months	Average	8.2%
Adolescent Mental Health and/or Depression Screening*		-	-
Colorectal Cancer Screening		-	-
Optimal Asthma Control - Adults		-	-
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		-	-
Optimal Vascular Care		-	-

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).