

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Above	68.2%
	Response at Six Months	Average	18.2%
	Remission at Six Months	Average	9.1%
	Follow-up at 12 Months	Average	61.4%
	Response at 12 Months	Average	20.5%
	Remission at 12 Months	Average	4.6%
Adult Depression	PHQ-9 Utilization*	Above	91.7%
	Follow-up at Six Months	Average	40.5%
	Response at Six Months	Average	13.2%
	Remission at Six Months	Average	5.9%
	Follow-up at 12 Months	Average	38.1%
	Response at 12 Months	Average	11.2%
	Remission at 12 Months	Average	4.4%
Adolescent Mental Health and/or Depression Screening*		Average	96.3%
Colorectal Cancer Screening		Below	55.2%
Optimal Asthma Control - Adults		Average	44.6%
Optimal Asthma Control - Children		Below	30.9%
Optimal Diabetes Care		Below	35.1%
Optimal Vascular Care		Average	47.3%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).