

Clinical Quality Measures

2022 measurement year

| Measure | | Rating | Rate |
|--|-------------------------|---------|-------|
| Adolescent Depression | PHQ-9 Utilization* | - | - |
| | Follow-up at Six Months | - | - |
| | Response at Six Months | - | - |
| | Remission at Six Months | - | - |
| | Follow-up at 12 Months | - | - |
| | Response at 12 Months | - | - |
| | Remission at 12 Months | - | - |
| Adult Depression | PHQ-9 Utilization* | Average | 77.1% |
| | Follow-up at Six Months | Below | 31.2% |
| | Response at Six Months | Average | 12.6% |
| | Remission at Six Months | Average | 8.4% |
| | Follow-up at 12 Months | Below | 30.6% |
| | Response at 12 Months | Average | 12.9% |
| | Remission at 12 Months | Average | 7.5% |
| Adolescent Mental Health and/or Depression Screening* | | Below | 83.6% |
| Colorectal Cancer Screening | | Above | 72.1% |
| Optimal Asthma Control - Adults | | Below | 33.4% |
| Optimal Asthma Control - Children | | Below | 18.8% |
| Optimal Diabetes Care | | Average | 40.0% |
| Optimal Vascular Care | | Below | 41.0% |

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).