

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	96.7%
	Follow-up at Six Months	Below	30.4%
	Response at Six Months	Average	10.9%
	Remission at Six Months	Average	6.5%
	Follow-up at 12 Months	Average	31.2%
	Response at 12 Months	Below	6.5%
	Remission at 12 Months	Average	3.6%
Adult Depression	PHQ-9 Utilization*	Below	66.4%
	Follow-up at Six Months	Below	32.6%
	Response at Six Months	Below	11.2%
	Remission at Six Months	Below	5.7%
	Follow-up at 12 Months	Below	35.5%
	Response at 12 Months	Below	11.3%
	Remission at 12 Months	Below	4.9%
Adolescent Mental Health and/or Depression Screening*		Above	97.6%
Colorectal Cancer Screening		Average	67.1%
Optimal Asthma Control - Adults		Below	37.2%
Optimal Asthma Control - Children		Below	41.0%
Optimal Diabetes Care		Below	36.9%
Optimal Vascular Care		Average	53.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).