

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	30.0%
	Response at Six Months	Average	10.0%
	Remission at Six Months	Average	3.3%
	Follow-up at 12 Months	Average	33.3%
	Response at 12 Months	Average	10.0%
	Remission at 12 Months	Average	6.7%
Adult Depression	PHQ-9 Utilization*	Average	79.4%
	Follow-up at Six Months	Average	49.0%
	Response at Six Months	Average	23.5%
	Remission at Six Months	Average	12.4%
	Follow-up at 12 Months	Average	52.3%
	Response at 12 Months	Average	24.2%
	Remission at 12 Months	Average	14.4%
Adolescent Mental Health and/or Depression Screening*		Above	98.8%
Colorectal Cancer Screening		Below	58.5%
Optimal Asthma Control - Adults		Below	32.8%
Optimal Asthma Control - Children		Below	33.7%
Optimal Diabetes Care		Average	41.7%
Optimal Vascular Care		Average	52.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).