

Clinical Quality Measures
 2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	30.4%
	Response at Six Months	Average	6.5%
	Remission at Six Months	Average	2.2%
	Follow-up at 12 Months	Average	37.0%
	Response at 12 Months	Average	19.6%
	Remission at 12 Months	Average	10.9%
Adult Depression	PHQ-9 Utilization*	Average	77.9%
	Follow-up at Six Months	Below	35.6%
	Response at Six Months	Below	10.8%
	Remission at Six Months	Average	6.5%
	Follow-up at 12 Months	Below	30.1%
	Response at 12 Months	Below	10.8%
	Remission at 12 Months	Average	6.0%
Adolescent Mental Health and/or Depression Screening*		Above	97.7%
Colorectal Cancer Screening		Average	66.7%
Optimal Asthma Control - Adults		Average	44.7%
Optimal Asthma Control - Children		Below	33.8%
Optimal Diabetes Care		Above	50.5%
Optimal Vascular Care		Average	56.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).