

Clinical Quality Measures
 2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	65.0%
	Follow-up at Six Months	Average	41.2%
	Response at Six Months	Average	10.5%
	Remission at Six Months	Below	1.8%
	Follow-up at 12 Months	Average	37.7%
	Response at 12 Months	Average	9.7%
	Remission at 12 Months	Average	4.4%
Adolescent Mental Health and/or Depression Screening*		Above	100.0%
Colorectal Cancer Screening		Average	67.4%
Optimal Asthma Control - Adults		Above	64.2%
Optimal Asthma Control - Children		Average	43.6%
Optimal Diabetes Care		Average	49.7%
Optimal Vascular Care		Average	58.3%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).