

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	84.8%
	Follow-up at Six Months	Average	46.9%
	Response at Six Months	Average	16.3%
	Remission at Six Months	Average	7.5%
	Follow-up at 12 Months	Average	42.2%
	Response at 12 Months	Average	17.0%
	Remission at 12 Months	Average	10.9%
Adult Depression	PHQ-9 Utilization*	Average	75.5%
	Follow-up at Six Months	Below	39.8%
	Response at Six Months	Below	13.5%
	Remission at Six Months	Below	7.1%
	Follow-up at 12 Months	Below	40.3%
	Response at 12 Months	Below	11.3%
	Remission at 12 Months	Below	6.4%
Adolescent Mental Health and/or Depression Screening*		Above	96.6%
Colorectal Cancer Screening		Above	73.6%
Optimal Asthma Control - Adults		Below	44.3%
Optimal Asthma Control - Children		Below	39.5%
Optimal Diabetes Care		Above	53.3%
Optimal Vascular Care		Above	66.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).