

Clinical Quality Measures

2022 measurement year

| Measure | | Rating | Rate |
|--|-------------------------|---------|-------|
| Adolescent Depression | PHQ-9 Utilization* | - | - |
| | Follow-up at Six Months | Average | 48.4% |
| | Response at Six Months | Average | 12.9% |
| | Remission at Six Months | Average | 6.5% |
| | Follow-up at 12 Months | Average | 35.5% |
| | Response at 12 Months | Average | 9.7% |
| | Remission at 12 Months | Average | 3.2% |
| Adult Depression | PHQ-9 Utilization* | Above | 93.4% |
| | Follow-up at Six Months | Average | 50.5% |
| | Response at Six Months | Average | 15.5% |
| | Remission at Six Months | Average | 9.6% |
| | Follow-up at 12 Months | Average | 47.8% |
| | Response at 12 Months | Average | 20.6% |
| | Remission at 12 Months | Average | 12.4% |
| Adolescent Mental Health and/or Depression Screening* | | Average | 92.4% |
| Colorectal Cancer Screening | | Average | 68.0% |
| Optimal Asthma Control - Adults | | Average | 40.7% |
| Optimal Asthma Control - Children | | Below | 24.3% |
| Optimal Diabetes Care | | Below | 39.0% |
| Optimal Vascular Care | | Below | 46.4% |

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).