

**Clinical Quality Measures**

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	77.1%
	Follow-up at Six Months	Average	38.6%
	Response at Six Months	Average	5.7%
	Remission at Six Months	Average	4.3%
	Follow-up at 12 Months	Average	35.7%
	Response at 12 Months	Average	8.6%
	Remission at 12 Months	Average	2.9%
Adult Depression	PHQ-9 Utilization*	Below	63.7%
	Follow-up at Six Months	Below	28.8%
	Response at Six Months	Below	10.5%
	Remission at Six Months	Below	3.8%
	Follow-up at 12 Months	Below	34.1%
	Response at 12 Months	Below	12.4%
	Remission at 12 Months	Below	6.2%
<b>Adolescent Mental Health and/or Depression Screening*</b>		Above	96.0%
<b>Colorectal Cancer Screening</b>		Below	61.0%
<b>Optimal Asthma Control - Adults</b>		Below	45.5%
<b>Optimal Asthma Control - Children</b>		Below	47.9%
<b>Optimal Diabetes Care</b>		Below	36.8%
<b>Optimal Vascular Care</b>		Average	56.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

**Ratings**

**Above:** Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).