

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	94.6%
	Follow-up at Six Months	Average	43.6%
	Response at Six Months	Average	11.5%
	Remission at Six Months	Average	5.1%
	Follow-up at 12 Months	Average	48.7%
	Response at 12 Months	Average	18.0%
	Remission at 12 Months	Average	11.5%
Adult Depression	PHQ-9 Utilization*	Above	83.2%
	Follow-up at Six Months	Below	38.4%
	Response at Six Months	Average	16.1%
	Remission at Six Months	Average	9.1%
	Follow-up at 12 Months	Below	38.2%
	Response at 12 Months	Average	14.9%
	Remission at 12 Months	Average	9.9%
Adolescent Mental Health and/or Depression Screening*		Above	98.9%
Colorectal Cancer Screening		Above	72.4%
Optimal Asthma Control - Adults		Above	62.5%
Optimal Asthma Control - Children		Average	55.3%
Optimal Diabetes Care		Above	57.1%
Optimal Vascular Care		Above	62.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).