

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	41.9%
	Response at Six Months	Average	16.3%
	Remission at Six Months	Average	7.0%
	Follow-up at 12 Months	Average	51.2%
	Response at 12 Months	Average	25.6%
	Remission at 12 Months	Average	11.6%
Adult Depression	PHQ-9 Utilization*	Average	77.1%
	Follow-up at Six Months	Average	41.0%
	Response at Six Months	Average	14.4%
	Remission at Six Months	Average	8.1%
	Follow-up at 12 Months	Below	36.2%
	Response at 12 Months	Average	14.0%
	Remission at 12 Months	Below	6.3%
Adolescent Mental Health and/or Depression Screening*		Above	99.7%
Colorectal Cancer Screening		Average	69.3%
Optimal Asthma Control - Adults		Average	54.9%
Optimal Asthma Control - Children		Average	61.7%
Optimal Diabetes Care		Average	50.8%
Optimal Vascular Care		Average	59.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).