

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	89.3%
	Follow-up at Six Months	Average	50.0%
	Response at Six Months	Average	13.5%
	Remission at Six Months	Average	7.7%
	Follow-up at 12 Months	Average	39.4%
	Response at 12 Months	Average	15.4%
	Remission at 12 Months	Average	6.7%
Adult Depression	PHQ-9 Utilization*	Below	65.4%
	Follow-up at Six Months	Below	41.2%
	Response at Six Months	Below	12.7%
	Remission at Six Months	Below	5.0%
	Follow-up at 12 Months	Average	40.0%
	Response at 12 Months	Below	12.0%
	Remission at 12 Months	Below	6.3%
Adolescent Mental Health and/or Depression Screening*		Above	94.9%
Colorectal Cancer Screening		Above	70.5%
Optimal Asthma Control - Adults		Average	47.1%
Optimal Asthma Control - Children		Below	30.5%
Optimal Diabetes Care		Average	43.2%
Optimal Vascular Care		Average	55.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).