

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	56.8%
	Response at Six Months	Average	18.2%
	Remission at Six Months	Average	6.8%
	Follow-up at 12 Months	Average	29.6%
	Response at 12 Months	Average	6.8%
	Remission at 12 Months	Average	6.8%
Adult Depression	PHQ-9 Utilization*	Above	87.0%
	Follow-up at Six Months	Average	41.7%
	Response at Six Months	Below	11.3%
	Remission at Six Months	Below	4.8%
	Follow-up at 12 Months	Average	45.2%
	Response at 12 Months	Average	16.1%
	Remission at 12 Months	Average	6.5%
Adolescent Mental Health and/or Depression Screening*		Above	98.4%
Colorectal Cancer Screening		Below	64.3%
Optimal Asthma Control - Adults		Average	60.4%
Optimal Asthma Control - Children		Average	51.3%
Optimal Diabetes Care		Average	53.3%
Optimal Vascular Care		Average	62.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).