

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	74.8%
	Follow-up at Six Months	Average	43.2%
	Response at Six Months	Below	9.5%
	Remission at Six Months	Below	4.5%
	Follow-up at 12 Months	Below	35.6%
	Response at 12 Months	Average	15.8%
	Remission at 12 Months	Average	8.6%
Adolescent Mental Health and/or Depression Screening*		-	-
Colorectal Cancer Screening		Above	80.1%
Optimal Asthma Control - Adults		Average	45.8%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Average	49.1%
Optimal Vascular Care		Average	58.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).