

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	53.9%
	Response at Six Months	Average	15.4%
	Remission at Six Months	Average	10.3%
	Follow-up at 12 Months	Average	51.3%
	Response at 12 Months	Average	25.6%
	Remission at 12 Months	Average	15.4%
Adult Depression	PHQ-9 Utilization*	Below	67.9%
	Follow-up at Six Months	Average	58.6%
	Response at Six Months	Average	21.9%
	Remission at Six Months	Average	11.7%
	Follow-up at 12 Months	Average	50.0%
	Response at 12 Months	Average	18.8%
	Remission at 12 Months	Average	4.7%
Adolescent Mental Health and/or Depression Screening*		Average	91.6%
Colorectal Cancer Screening		Below	63.9%
Optimal Asthma Control - Adults		Above	62.7%
Optimal Asthma Control - Children		Above	73.7%
Optimal Diabetes Care		Below	38.5%
Optimal Vascular Care		Average	55.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).