

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	76.9%
	Follow-up at Six Months	Average	54.0%
	Response at Six Months	Average	20.0%
	Remission at Six Months	Average	16.0%
	Follow-up at 12 Months	Average	56.0%
	Response at 12 Months	Average	26.0%
	Remission at 12 Months	Average	12.0%
Adolescent Mental Health and/or Depression Screening*		Average	94.1%
Colorectal Cancer Screening		Average	72.4%
Optimal Asthma Control - Adults		Average	54.6%
Optimal Asthma Control - Children		Average	47.1%
Optimal Diabetes Care		Average	53.7%
Optimal Vascular Care		Average	60.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).