

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	92.9%
	Follow-up at Six Months	Average	49.1%
	Response at Six Months	Average	12.7%
	Remission at Six Months	Average	4.6%
	Follow-up at 12 Months	Average	45.5%
	Response at 12 Months	Average	10.0%
	Remission at 12 Months	Average	3.6%
Adult Depression	PHQ-9 Utilization*	Average	78.3%
	Follow-up at Six Months	Below	40.5%
	Response at Six Months	Below	12.1%
	Remission at Six Months	Below	5.8%
	Follow-up at 12 Months	Below	39.1%
	Response at 12 Months	Average	14.0%
	Remission at 12 Months	Average	8.1%
Adolescent Mental Health and/or Depression Screening*		Above	94.8%
Colorectal Cancer Screening		Above	72.2%
Optimal Asthma Control - Adults		Below	46.7%
Optimal Asthma Control - Children		Below	36.6%
Optimal Diabetes Care		Average	45.5%
Optimal Vascular Care		Average	57.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).