

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	95.7%
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Below	32.1%
	Response at Six Months	Average	13.1%
	Remission at Six Months	Average	6.0%
	Follow-up at 12 Months	Average	33.3%
	Response at 12 Months	Average	9.5%
	Remission at 12 Months	Average	7.1%
Adolescent Mental Health and/or Depression Screening*		Below	78.1%
Colorectal Cancer Screening		Below	54.3%
Optimal Asthma Control - Adults		Above	69.0%
Optimal Asthma Control - Children		Average	51.6%
Optimal Diabetes Care		Average	43.2%
Optimal Vascular Care		Average	49.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).