

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	58.5%
	Response at Six Months	Average	22.6%
	Remission at Six Months	Above	18.9%
	Follow-up at 12 Months	Average	49.1%
	Response at 12 Months	Average	15.1%
	Remission at 12 Months	Average	5.7%
Adult Depression	PHQ-9 Utilization*	Above	82.3%
	Follow-up at Six Months	Average	41.7%
	Response at Six Months	Average	14.0%
	Remission at Six Months	Average	7.8%
	Follow-up at 12 Months	Average	38.4%
	Response at 12 Months	Average	14.4%
	Remission at 12 Months	Average	7.4%
Adolescent Mental Health and/or Depression Screening*		Above	98.1%
Colorectal Cancer Screening		Average	66.5%
Optimal Asthma Control - Adults		Average	48.2%
Optimal Asthma Control - Children		Average	48.4%
Optimal Diabetes Care		Above	50.7%
Optimal Vascular Care		Above	63.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).