

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	54.8%
	Response at Six Months	Average	16.1%
	Remission at Six Months	Average	6.5%
	Follow-up at 12 Months	Average	48.4%
	Response at 12 Months	Average	19.4%
	Remission at 12 Months	Average	16.1%
Adult Depression	PHQ-9 Utilization*	Average	79.1%
	Follow-up at Six Months	Below	36.9%
	Response at Six Months	Average	12.1%
	Remission at Six Months	Average	8.7%
	Follow-up at 12 Months	Below	34.9%
	Response at 12 Months	Below	10.1%
	Remission at 12 Months	Average	6.4%
Adolescent Mental Health and/or Depression Screening*		Average	93.2%
Colorectal Cancer Screening		Above	74.9%
Optimal Asthma Control - Adults		Average	45.6%
Optimal Asthma Control - Children		Average	50.0%
Optimal Diabetes Care		Average	49.8%
Optimal Vascular Care		Average	62.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).