

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	32.4%
	Response at Six Months	Average	13.5%
	Remission at Six Months	Average	8.1%
	Follow-up at 12 Months	Average	29.7%
	Response at 12 Months	Average	8.1%
	Remission at 12 Months	Average	0.0%
Adult Depression	PHQ-9 Utilization*	Average	77.4%
	Follow-up at Six Months	Below	39.1%
	Response at Six Months	Average	17.3%
	Remission at Six Months	Average	8.4%
	Follow-up at 12 Months	Below	32.4%
	Response at 12 Months	Below	9.3%
	Remission at 12 Months	Below	5.3%
Adolescent Mental Health and/or Depression Screening*		Above	98.0%
Colorectal Cancer Screening		Above	72.0%
Optimal Asthma Control - Adults		Average	50.2%
Optimal Asthma Control - Children		Average	50.0%
Optimal Diabetes Care		Average	49.9%
Optimal Vascular Care		Average	58.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).