

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	94.2%
	Follow-up at Six Months	Average	48.2%
	Response at Six Months	Average	12.5%
	Remission at Six Months	Average	3.6%
	Follow-up at 12 Months	Average	39.3%
	Response at 12 Months	Average	16.1%
	Remission at 12 Months	Average	7.1%
Adult Depression	PHQ-9 Utilization*	Above	82.4%
	Follow-up at Six Months	Above	57.1%
	Response at Six Months	Average	19.8%
	Remission at Six Months	Average	10.3%
	Follow-up at 12 Months	Above	54.1%
	Response at 12 Months	Average	20.7%
	Remission at 12 Months	Average	10.3%
Adolescent Mental Health and/or Depression Screening*		Below	85.0%
Colorectal Cancer Screening		Below	67.5%
Optimal Asthma Control - Adults		Above	58.9%
Optimal Asthma Control - Children		Average	56.4%
Optimal Diabetes Care		Above	51.1%
Optimal Vascular Care		Average	57.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).