

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Average	61.5%
	Response at Six Months	Average	20.5%
	Remission at Six Months	Average	7.7%
	Follow-up at 12 Months	Average	51.3%
	Response at 12 Months	Average	10.3%
	Remission at 12 Months	Average	7.7%
Adult Depression	PHQ-9 Utilization*	Above	99.1%
	Follow-up at Six Months	Above	60.4%
	Response at Six Months	Average	22.9%
	Remission at Six Months	Average	14.2%
	Follow-up at 12 Months	Average	41.1%
	Response at 12 Months	Average	13.5%
	Remission at 12 Months	Average	8.4%
Adolescent Mental Health and/or Depression Screening*		Above	98.7%
Colorectal Cancer Screening		Above	78.5%
Optimal Asthma Control - Adults		Above	65.7%
Optimal Asthma Control - Children		Average	62.5%
Optimal Diabetes Care		Average	47.9%
Optimal Vascular Care		Average	57.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).