

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	46.7%
	Response at Six Months	Average	20.0%
	Remission at Six Months	Average	13.3%
	Follow-up at 12 Months	Average	23.3%
	Response at 12 Months	Average	13.3%
	Remission at 12 Months	Average	3.3%
Adult Depression	PHQ-9 Utilization*	Above	97.4%
	Follow-up at Six Months	Average	52.7%
	Response at Six Months	Average	22.3%
	Remission at Six Months	Average	9.6%
	Follow-up at 12 Months	Average	38.2%
	Response at 12 Months	Average	13.2%
	Remission at 12 Months	Average	7.7%
Adolescent Mental Health and/or Depression Screening*		Above	96.0%
Colorectal Cancer Screening		Average	69.0%
Optimal Asthma Control - Adults		Average	47.1%
Optimal Asthma Control - Children		Average	49.3%
Optimal Diabetes Care		Average	41.9%
Optimal Vascular Care		Average	48.3%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).