

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	75.4%
	Follow-up at Six Months	Below	32.6%
	Response at Six Months	Below	9.8%
	Remission at Six Months	Below	5.4%
	Follow-up at 12 Months	Below	32.1%
	Response at 12 Months	Below	10.9%
	Remission at 12 Months	Below	5.4%
Adolescent Mental Health and/or Depression Screening*		Below	88.4%
Colorectal Cancer Screening		Above	71.0%
Optimal Asthma Control - Adults		Below	41.2%
Optimal Asthma Control - Children		Below	31.3%
Optimal Diabetes Care		Below	39.5%
Optimal Vascular Care		Below	38.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).