

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	94.1%
	Follow-up at Six Months	Average	60.0%
	Response at Six Months	Average	23.3%
	Remission at Six Months	Average	8.3%
	Follow-up at 12 Months	Average	51.7%
	Response at 12 Months	Average	18.3%
	Remission at 12 Months	Average	10.0%
Adult Depression	PHQ-9 Utilization*	Above	84.6%
	Follow-up at Six Months	Above	64.2%
	Response at Six Months	Above	27.6%
	Remission at Six Months	Above	17.4%
	Follow-up at 12 Months	Above	68.2%
	Response at 12 Months	Above	31.4%
	Remission at 12 Months	Above	20.1%
Adolescent Mental Health and/or Depression Screening*		Above	99.1%
Colorectal Cancer Screening		Above	76.7%
Optimal Asthma Control - Adults		Below	38.8%
Optimal Asthma Control - Children		Average	55.1%
Optimal Diabetes Care		Above	53.9%
Optimal Vascular Care		Average	62.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).