

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	62.2%
	Follow-up at Six Months	Above	76.0%
	Response at Six Months	Above	38.5%
	Remission at Six Months	Above	26.0%
	Follow-up at 12 Months	Above	63.5%
	Response at 12 Months	Above	34.4%
	Remission at 12 Months	Above	20.8%
Adolescent Mental Health and/or Depression Screening*		Average	91.4%
Colorectal Cancer Screening		Below	63.0%
Optimal Asthma Control - Adults		Average	48.9%
Optimal Asthma Control - Children		Average	58.1%
Optimal Diabetes Care		Average	40.6%
Optimal Vascular Care		Average	53.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).