

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	67.0%
	Follow-up at Six Months	Above	82.4%
	Response at Six Months	Above	38.2%
	Remission at Six Months	Above	22.1%
	Follow-up at 12 Months	Above	70.6%
	Response at 12 Months	Above	32.4%
	Remission at 12 Months	Above	20.1%
Adolescent Mental Health and/or Depression Screening*		Above	97.4%
Colorectal Cancer Screening		Average	65.6%
Optimal Asthma Control - Adults		Above	61.7%
Optimal Asthma Control - Children		Average	60.5%
Optimal Diabetes Care		Above	49.6%
Optimal Vascular Care		Above	60.3%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).