

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	85.7%
	Follow-up at Six Months	Average	59.2%
	Response at Six Months	Average	22.5%
	Remission at Six Months	Average	8.2%
	Follow-up at 12 Months	Average	49.0%
	Response at 12 Months	Average	20.4%
	Remission at 12 Months	Average	10.2%
Adult Depression	PHQ-9 Utilization*	Below	68.5%
	Follow-up at Six Months	Above	81.1%
	Response at Six Months	Above	40.0%
	Remission at Six Months	Above	27.7%
	Follow-up at 12 Months	Above	73.0%
	Response at 12 Months	Above	35.5%
	Remission at 12 Months	Above	24.5%
Adolescent Mental Health and/or Depression Screening*		Above	97.8%
Colorectal Cancer Screening		Above	69.5%
Optimal Asthma Control - Adults		Above	57.8%
Optimal Asthma Control - Children		Average	51.6%
Optimal Diabetes Care		Average	44.9%
Optimal Vascular Care		Average	53.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).