

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	55.8%
	Response at Six Months	Average	16.3%
	Remission at Six Months	Average	11.6%
	Follow-up at 12 Months	Average	41.9%
	Response at 12 Months	Average	14.0%
	Remission at 12 Months	Average	2.3%
Adult Depression	PHQ-9 Utilization*	Below	64.8%
	Follow-up at Six Months	Above	81.4%
	Response at Six Months	Above	41.4%
	Remission at Six Months	Above	26.0%
	Follow-up at 12 Months	Above	72.4%
	Response at 12 Months	Above	37.2%
	Remission at 12 Months	Above	23.7%
Adolescent Mental Health and/or Depression Screening*		Average	95.6%
Colorectal Cancer Screening		Above	73.8%
Optimal Asthma Control - Adults		Above	72.4%
Optimal Asthma Control - Children		Above	76.3%
Optimal Diabetes Care		Average	49.7%
Optimal Vascular Care		Average	60.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).