

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	94.7%
	Follow-up at Six Months	Above	66.0%
	Response at Six Months	Above	23.4%
	Remission at Six Months	Average	9.0%
	Follow-up at 12 Months	Above	58.5%
	Response at 12 Months	Above	24.5%
	Remission at 12 Months	Average	11.2%
Adult Depression	PHQ-9 Utilization*	Below	41.4%
	Follow-up at Six Months	Above	76.3%
	Response at Six Months	Above	32.2%
	Remission at Six Months	Above	20.1%
	Follow-up at 12 Months	Above	71.7%
	Response at 12 Months	Above	27.6%
	Remission at 12 Months	Average	14.0%
Adolescent Mental Health and/or Depression Screening*		Above	98.2%
Colorectal Cancer Screening		Above	72.5%
Optimal Asthma Control - Adults		Above	58.7%
Optimal Asthma Control - Children		Above	75.8%
Optimal Diabetes Care		Average	48.0%
Optimal Vascular Care		Above	63.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).