

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	94.0%
	Follow-up at Six Months	Average	63.3%
	Response at Six Months	Average	18.4%
	Remission at Six Months	Average	6.1%
	Follow-up at 12 Months	Average	49.0%
	Response at 12 Months	Average	14.3%
	Remission at 12 Months	Average	8.2%
Adult Depression	PHQ-9 Utilization*	Below	64.8%
	Follow-up at Six Months	Above	78.3%
	Response at Six Months	Above	34.8%
	Remission at Six Months	Above	17.4%
	Follow-up at 12 Months	Above	69.1%
	Response at 12 Months	Above	31.2%
	Remission at 12 Months	Above	17.6%
Adolescent Mental Health and/or Depression Screening*		Above	98.7%
Colorectal Cancer Screening		Above	73.9%
Optimal Asthma Control - Adults		Above	63.3%
Optimal Asthma Control - Children		Average	55.3%
Optimal Diabetes Care		Above	53.5%
Optimal Vascular Care		Above	64.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).